

PAROXETINA – propunere protocol

1. Denumire stiintifica:
Paroxetinum
2. Clasa de medicamente:
Antidepresive (ATC: N06AB05)
3. Forme farmaceutice (ANMDM):
 - comprimate 10 mg, 20 mg, 30 mg, 40 mg
 - comprimate filmate 20 mg, 40 mg
4. Profil farmacologic : ISRS (inhibitori selectivi ai recaptarii de serotonina) (2 pag. 415)
5. Farmacocinetica :
 - timp de injumatatire de aproximativ 24 de ore, metabolit inactiv, metabolizare hepatica, inhiba CYP 2D6
6. Mecanism de actiune: blocheaza pompa de recaptare a serotoninei, crescand neurotransmisia serotoninergica. Poate avea activitate de blocant slab al recaptarii de noradrenalina. Are si actiune anticolinergica slaba (1 pag 415, 2 pag 298)
Dozare : 20-60mg
7. Indicatii principale (ANMDM, EMA, FDA) (8, 9, 10):
 - Episod depresiv major
 - Tulburare obsesiv-compulsiva
 - Tulburare de panica cu sau fara agorafobie
 - Tulburare de anxietate sociala/Fobie sociala
 - Tulburare de anxietate generalizata
 - Tulburare de stres post-traumatic
8. Alte indicatii:
 - Tulburare disforica premenstruala (FDA, 2 pag 415, 6 pag 1014)
 - Simptome vasomotorii moderat-severe asociate menopauzei (FDA: cps 7,5mg)
 - Episod depresiv din TAB in combinatie cu timostabilizatoare (5)
 - Depresie si Comorbiditati somatice (indicatii ISRS) : Boli cardiovasculare – Boala cardiaca ischemica (11, 12, 13, 14) , AVC (11, 13); Afectiuni oncologice (12, 13); Diabet zaharat (11, 12, 13); Epilepsie (11, 12); Demente (12, 13); Infectie HIV (11, 12).
 - Tulburarea depresiva persistenta (Distimia si Depresia cronica): ISRS (13, 14, 15, 16).
 - Schizofrenie (indicatii antidepresive): - depresia majora comorbida, depresia post-psihotica, TOC (12, 17, 18, 19, 20, 21, 22); simptome depresive severe, care determina disconfort semnificativ sau interfereaza cu functionarea (17, 22).
9. Efecte secundare : greata, senzatie de gura uscata, ameteli, tremor, constipatie sau diaree, disfunctii sexuale , aplatizare afectiva , lentoare cognitiva , insomnie sau sedare, crestere in greutate , akatisie (2 pag 416, 6 pag 583, 6 pag 1018)
Intreruperea paroxetinei nu se va face brusc din cauza sindromului de discontinuitate. Simptomele de intrerupere (ameteala, greata, crampe, diaforeza, parestezii, disestezii) tind sa fie mai severe sau mai frecvente decat in cazul

celorlalte ISRS-uri. La cei cu reactii severe va intrerupe foarte lent (saptamani /luni) sau se va asocia cu un ISRS cu timp lung de injumatare (ex, fluoxetina) . Mecanismul implica inhibitia propriului metabolism la intreruperea brusca, cu rata de scadere brusca a concentratiei plasmatice la care se adauga readaptarea receptorilor colinergici dupa blocarea lor prelungita.

10. Supradoza:

- toxicitate scazuta in supradoza; sindrom serotoninergic, sedare, voma, midriaza, tulburari de ritm cardiac. Forte rar s-au raportat cazuri fatale, in combinatie cu alte medicamente/substante psihoactive (2 pag 418) (rcp)

11. Utilizare la grupe de pacienti cu risc crescut:

- Afectare renala: se recomanda utilizarea unor doze mai scazute, dar poate fi data si pacientilor dializati (2 pag 419, 3 pag 583)
- Afectare hepatica: se utilizeaza doze scazute (10 mg in insuficienta hepatica severa) (2 pag 419, 3 pag 583)
- Afectare cardiaca: prezinta siguranta la aceasta categorie de pacienti (2 pag 419)
- Varstnici: se utilizeaza doze reduse (maxim 40 mg).
- Sarcina: categorie de risc D, a fost asociata in principal cu malformatii cardiace la fat, desi unele studii au infirmat acest lucru. Au fost raportate complicatii neonatale, inclusiv detresa respiratorie (posibil si datorita sindromului de intrerupere), hipertensiune pulmonara la nou-nascut. Se pare ca este mai putin sigura decat alte ISRS-uri la aceasta categorie. Trebuie evaluate riscurile tratamentului pentru fat fata de riscurile absentei tratamentului atat pentru mama cat si pentru copil .(1 pag 361, 2 pag 420, 3 pag 547)
- Alaptare: cantitati minime de medicament trec in laptele matern. Se considera ca poate fi dat in alaptare, dar daca sugarul devine iritabil sau sedat, se recomanda trecerea la alimentatie artificiala , intreruperea tratamentului sau utilizarea unui antidepresiv cu un profil mai sigur. In toate cazurile se cantaresc riscurile tratamentului cu beneficiile tratarii depresiei atat pentru mama cat si pentru copil (2 pag 420, 3 pag 563, 4 pag 83).
- Epilepsie : se recomanda precautie (risc scazut de convulsii). Nu afecteaza farmacocinetica/farmacodinamia anticonvulsivantelor (rcp).

12. Interactiuni medicamentoase :

- poate creste riscul de hemoragii mai ales la pacientii cu anticoagulante/agenti antiplachetari. Interfereaza cu analgezicele opioide . Nu se administreaza cu pimozid (creste concentratia plasmatica a pimozidului , iar acesta prelungeste intervalul QTc) (2 pag 419, 6 pag 1014). Nu necesita ajustarea dozei cand se administreaza impreuna cu alte inductoare enzimatice (carbamazepina, rifampicina, fenobarbital, fenitoina) (rcp)
 - creste nivelul plasmatic al unor antipsihotice, beta-blocante, al tioridazinei (2 pag 418-419, 3 pag 332 ,4 pag 83)
- Contraindicatii : tratament cu IMAO, tioridazina, pimozid
- Alte interactiuni : nu se recomanda consumul de alcool

Referinte:

1. Stahl SM., Stahl's Essential Psychopharmacology: Neuroscientific Basis and Practical Application 4th edition. New York: Cambridge University Press 2013
2. Stahl MS- Psihofarmacologie –Ghidul prescriptorului ed. a 3-a revizuita 2012
3. Maudsley Prescribing Guidelines in Psychiatry 12th ed.
4. BAP Fundamentals of Clinical Psychopharmacology 4th ed. 2016
5. American Pszchiatric Association Practice Guideline for the Treatment of Patients With Bipolar Disorder Second Edition, 2002. Hirschfeld RMA: Guideline Watch: Practice Guideline for the Treatment of Patients With Bipolar Disorder. Arlington, VA: American Psychiatric Association.
6. Kaplan synopsis of Psychiatry 11th Ed
7. http://www.ema.europa.eu/docs/en_GB/document_library/Referrals_document/Paroxetine_31/WC500012852.pdf
8. ANMDM - Nomenclatorul medicamentelor pentru uz uman
https://www.anm.ro/_/RCP/rcp_4180_18.01.12.pdf
9. EMA European Medicines Agency
http://www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/referrals/Paroxetine/human_referral_000131.jsp&mid=WC0b01ac05805c516f
10. FDA U.S. Food and Drug Administration
https://www.accessdata.fda.gov/drugsatfda_docs/label/2012/020031s067,020710s031.pdf
https://www.accessdata.fda.gov/drugsatfda_docs/label/2012/020936s045lbl.pdf
https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/204516s005lbl.pdf
11. American Psychiatric Association, Practice Guideline for the Treatment of Patients with Major Depressive Disorder 3rd ed November 2010.
12. Bon Usage des Medicaments Antidepresseurs dans le Traitement des Troubles Depressifs et des Troubles Anxieux de L'Adulte, Agence Française de Sécurité Sanitaire des Produits de Santé, octobre 2006, www.afssaps.sante.fr
13. German Society for Psychiatry and Psychotherapy, Psychosomatics and Neurology, S3 Guideline Unipolar Depression, 2015/ DGPPN, BÄK, KBV, AWMF, S3-Leitlinie/Nationale VersorgungsLeitlinie Unipolare Depression, Langfassung 2. Auflage, Version 5, 2015.
14. Mahli G.S., Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for mood disorders, Australian and New Zealand Journal of Psychiatry 2015, Vol. 49(12) 1-185.
15. Canadian Network for Mood and Anxiety Treatments (CANMAT) Guidelines for the management of adults with major depressive disorder, The Canadian Journal of Psychiatry / La Revue Canadienne de Psychiatrie 2016, Vol. 61(9) 504-505.
16. Bauer M. et al., World Federation of Societies of Biological Psychiatry (WFSBP) Guidelines for Biological Treatment of Unipolar Depressive Disorders in Primary Care, The World Journal of Biological Psychiatry, 2007; 8(2): 67-104.
17. Lehman A.F. et al., American Psychiatric Association, Practice Guideline for the Treatment of Patients with Schizophrenia Second Edition, 2004.
18. Remington G. et al., Canadian Schizophrenia Guidelines, Guidelines for the Pharmacotherapy of Schizophrenia in Adults, The Canadian Journal of Psychiatry / La Revue Canadienne de Psychiatrie 2017, Vol. 62(9) 604-616.

19. Hasan A. et al., World Federation of Societies of Biological Psychiatry (WFSBP) Guidelines for Biological Treatment of Schizophrenia. Part 3: Update 2015 Management of special circumstances: Depression, Suicidality, substance use disorders and pregnancy and lactation, The World Journal of Biological Psychiatry, 2015; 16: 142–170.
20. Galletly C. et al., Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for the management of schizophrenia and related disorders, Australian and New Zealand Journal of Psychiatry 2016, Vol. 50(5) 1-117.
21. Scottish Intercollegiate Guidelines Network, Management of schizophrenia, A national clinical guideline, March 2013.
22. Clinical Practice Guidelines In the NHS Ministry Of Health And Consumer Affairs, Clinical Practice Guideline for Schizophrenia and Incipient Psychotic Disorder, Published by: Catalan Agency for Health Technology Assessment and Research , 1st ed., 2009.